

DTIC FILE COPY

The silvered leaf monkey (Presbytis cristata) as a model
for human bancroftian filariasis

(2)


James R. Campbell, Harijani A. Marwoto, Soekartono Tirtokusumo,
Sofyan Masbar, Jan T. Rusch, Purnomo, and Bambang Trenggono

REPORT NO.

TR - 1044

AD-A201 887

DTIC
ELECTE
NOV 03 1988
S D
CVD



**UNITED STATES NAVAL
MEDICAL RESEARCH UNIT NO. TWO**

APO SAN FRANCISCO, CALIFORNIA 96328

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

88 1031 I16

ADMINISTRATIVE INFORMATION

C.G. HAYES, Ph.D.
Scientific Director

F.P. PALEOLOGO
Officer in Charge
Jakarta Detachment

This study was supported through funds provided by the Naval Medical Research and Development Command, Navy Department, for Work Unit 3M162770A870 AN314, and by the Indonesian Ministry of Health.

Distribution of this document is unlimited.

J.C. COOLBAUGH
CAPT MSC USN
Commanding Officer



Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-120	

The Silvered Leaf Monkey (*Presbytis cristata*) as a Model for Human Bancroftian Filariasis.

James R Campbell, Harijani A Marwoto, Soekartono Tirtokusumo, Sofyan Masbar, Jan T Rusch, Purnomo, and Bambang Trenggono

The silvered leaf monkey, *Presbytis cristata*, (Ceropithecidae:Colobinae) is the first animal other than man that has been shown capable of readily supporting the complete development of the human filarial parasite *Wuchereria bancrofti* (1,2). Attempts to infect other primates (*Macaca*) resulted in patent infections, but often required prior immunosuppression of the host. In all cases microfilarial densities were low, and periodicity could not be determined (3). Attempts to produce patent infections in smaller laboratory animals have been uniformly unsuccessful (4,5). Since *P. cristata* readily develop patent *W. bancrofti* infections without prior host immunosuppression, and because this particular primate species is abundant in Indonesia, it was chosen as a model for detailed studies of bancroftian filariasis.

To be useful as an experimental model for studying bancroftian filariasis, wild caught *P. cristata* first must be thoroughly characterized by physical, hematological and parasitological parameters. This study reports such baseline data obtained from a large sample of *P. cristata*.

A total of 169 *P. cristata* from Lampung, South Sumatra and Banten, West Java, Indonesia, were used in the study. The animals were caught in the wild and brought to a primate research facility in Jakarta, Indonesia, within 2-12 days after capture. Within 24 hours' after arrival, each animal was restrained with an intramuscular injection of ketamine HCl (Vetalar, Parke-Davis, Morris Plains, NJ, 07950), 10 mg/kg body weight, and given the following examinations: (a) Thick and thin blood smears for malaria examination and a 30 μ l thick smear for filaria examination were prepared and stained with Giemsa. In addition 1cc of heparinized blood drawn between 2200-0200 hours was filtered through a 3 μ m nucleopore filter, which was stained and examined for microfilariae. Another thin blood smear was stained with Wright-Giemsa and used for differential count. Hematocrit was determined by the micro-hematocrit method, hemoglobin was determined by the acid-hematin test and

total white blood cell counts were made using a Neubauer hemacytometer.

(b) Physical examinations were conducted to determine sex, weight, dentition and general physical condition, and any current or old injuries were noted.

(c) Each animal was given a tuberculosis skin test in the right eyelid consisting of an intradermal injection of 0.1cc of tuberculin (Mammalian, human isolates, Burroughs Wellcome, Kansas City, MO, 64161). Skin test were repeated at 2 weeks and at 4 weeks after arrival.

(d) Feces were collected, placed in formalin and in polyvinyl alcohol (PVA) and examined for intestinal parasites. Rectal swabs were taken, placed in Amies transport media and examined within 12 hours for bacterial pathogens.

(e) Animals were placed on white paper and their entire body was thoroughly brushed with a stiff brush to dislodge ectoparasites. Further careful examinations of the skin and the ears were conducted to locate ectoparasites.

Of the 169 animals in the study, 137 (81%) were female and 32 (19%) were male. Female monkeys averaged 4.5 ± 1.3 kg (range 1.6 - 6.7 kg) and males averaged 4.5 ± 1.8 kg (range 1.6 - 8.2 kg) in body weight. Blood parameters determined for these animals are summarized in Table 1.

Examination of the monkey's blood and feces revealed a variety of naturally acquired parasitic infections, as shown in Table 2. Amoebae found in stools included *E. polecki* and *E. hartmanni*, but no *E. histolytica* was found. In addition, in several animals *Oesophagostomum* sp. was found at necropsy in nodules on the serosal surface of the large and small intestines.

Bacteriological findings revealed a very low prevalence of intestinal pathogens, as shown in Table 3. All animals were negative on three successive skin tests for tuberculosis. No ectoparasites were found on any of the animals.

Table 1 Baseline hematological parameters in wild caught *Presbytis cristata*

parameter	n	X \pm S.D.	range
Hemoglobin (g dl)	169	9.6 ± 1.4	4.0 - 15.0
Hematocrit (%)	169	35 ± 4.6	18 - 48
WBC mm ³	164	8155 ± 3347	2100 - 18,600
Neutrophils (%)	93	50 ± 21	4 - 94
Lymphocytes (%)	93	48 ± 21	6 - 96
Eosinophils (%)	93	1 ± 5	0 - 13
Monocytes (%)	93	< 1	0 - 3
Ba-ophils (%)	93	< 1	0 - 2

From the U.S. Naval Medical Research Unit No. 2, Jakarta Detachment, APO San Francisco, CA 96356 (Campbell, Tirtokusumo, Masbar, Rusch, Purnomo, Trenggono) and the Indonesian Ministry of Health Research and Development, Jakarta, Indonesia (Marwoto)

Table 2 Naturally acquired parasitic infections in *Presbytis cristata*

species	number positive	number examined (%)
<i>Plasmodium knowlesi</i> (blood stages)	25/149	(16.8)
<i>Brugia malayi</i> (microfilariae)	7/132	(5.3)
<i>Trichuris trichiura</i> (ova)	123/158	(77.8)
<i>Strongyloides</i> sp. (larvae)	30/158	(19.0)
Ancylostomatids (ova)	13/158	(8.2)
<i>Entamoeba</i> sp. (cysts)	49/158	(31.0)
<i>Giardia lamblia</i> (cysts)	3/158	(1.9)

Table 3 Bacteriological culture results from rectal swabs of wild caught *Presbytis cristata*

species	number positive	number examined (%)
<i>Salmonella</i> group B ^b	8/180	(4.4)
<i>Salmonella</i> group E ^c	2/180	(1.1)
<i>Vibrio cholerae</i> non-O1 ^d	1/180	(0.6)

^aData represent the number of animals with rectal swabs culture-positive for organisms, versus total number of animals examined. Data are included from 11 additional animals that died before complete examinations could be performed.

^bSeven of these isolates were tetracycline resistant *in vitro*, one was tetracycline and ampicillin resistant.

^cBoth isolates were tetracycline resistant.

^dIsolate was colistin sulfate resistant.




Figure 1 Dentition pattern versus bodyweight in wild caught *Presbytis cristata* (n = 169). Solid lines = upper jaw dashed lines = lower jaw.

However lung mites, *Pneumonyssus simicola*, were later found in pathological specimens from a number of animals.

Approximately 25% of the animals arrived with fresh minor lacerations, for which appropriate treatment was given, and approximately 15% displayed scars, healed fractures or evidence of other old injuries. Two females arrived pregnant but both aborted within 24 hours after arrival. A comparison of dentition patterns with body weight upon arrival is shown in Figure 1. Sex ratio and mean body weight of the animals collected for use in this study were similar to values reported by Walker, et al. (6) for *P. cristata* in Malaysia.

Hematological parameters of the animals used in this study displayed a wide range of values. This variance probably reflects the difference in state of health of individual animals upon arrival. White blood cell differential counts were similar to those reported by Walker, et al. (6), although the latter were determined for animals after 3 months in captivity. Erythrocyte associated values for *P. cristata* gradually increase with time in captivity, sug-

gesting improvement of an anemic condition common among animals in the wild (6).

P. cristata are arboreal animals that avoid contact with humans, however the monkeys in this study were found to be naturally infected with several species of parasitic organisms common to man. In most cases, the short time between capture of the animals in the field and examination in the laboratory precluded acquisition of these infections from human animal trappers and handlers. Parasitological results from this study differed from those of Palmieri, et al. (7), however the latter study reported results from animals captured in a different geographical region of Indonesia. Arambulo, et al. (8) also reported high prevalence rates of *T. trichiura* and *Strongyloides* among wild caught *P. cristata* in Malaysia.

Culture of rectal swabs obtained on arrival detected pathogenic microorganisms in only a few animals. This could reflect a less than optimal isolation method rather than a lack of pathogens. However, with the increased human contact that accompanied captivity, the monkeys began experiencing frequent episodes of diarrhea. This was a serious problem that required constant vigilance and aggressive treatment with antibiotics and rehydration. A common and serious finding among new animals was respiratory infection, which if not aggressively treated was often fatal. A common etiology for this illness was not established, however the animals frequently responded well to trimethoprim-sulfamethoxazole or penicillin-streptomycin therapy. In many cases, it appeared probable that the animals' illness was exacerbated by the stress of captivity and handling.

Although procedures for locating ectoparasites on the animals were not optimal, the complete absence of findings suggests that in the wild state, the *P. cristata* used in our study did not carry heavy burdens of ectoparasites. However, lung mites were found at inflammatory foci in pathologic specimens from several monkeys, and may predispose their hosts to respiratory infections.

New animals presented with various minor injuries, particularly lacerations of the scalp, which we observed to result from stress behavior of striking the head against the cage and bites received from other captive monkeys. Figure 1 shows mean body weights at which various dental patterns were observed. Although these data cannot establish a clear correlation between dental pattern and age of the animal, as has been established for *Papio* sp. (9), they represent the first step in producing such a correlation for the better understanding of this important yet fragile animal.

Acknowledgements

This study was supported by work unit no. 3M162770A870.AN.314 from the Naval Medical Research and Development Command, Bethesda, Maryland and the Indonesian Ministry of Health.

We thank Mr. Chuck Darsono and his staff at the Cengkareng Primate Laboratory for providing facilities and services. We acknowledge Dr. I. Koiman, Director of the Biomedical

Research Institute of the Indonesian Ministry of Health Research and Development, for his support and encouragement.

The views of the authors do not purport to reflect the positions of the U.S. Department of the Navy, the naval service at large, or the Indonesian Ministry of Health.

References

1. Harinasuta C, Sucharit S, Choocote W. The susceptibility of leaf monkeys to bancroftian filariasis in Thailand. *SE Asian J Trop Med Pub Hlth* 1981; 12:581-89.
2. Palmieri JR, Connor DH, Purnomo, *et al.* Experimental infection of *Wuchereria bancrofti* in the silvered leaf monkey *Presbytis cristatus* Eschscholtz, 1821. *J Helminthol* 1982; 56:243-45.
3. Cross JH, Partono F, Hsu M-Y K, *et al.* Experimental transmission of *Wuchereria bancrofti* to monkeys. *Am J Trop Med Hyg* 1979; 28:56-66.
4. Dissanaikie AS. Developing larvae of *Wuchereria bancrofti* in the pulmonary vessels of splenectomized hamsters. *SE Asian J Trop Med Pub Hlth* 1974; 5:139-40.
5. Suswillo RR, Nelson GS, Muller R, *et al.* Attempts to infect jirds (*Meriones unguiculatus*) with *Wuchereria bancrofti*, *Onchocerca volvulus*, *Loa loa* and *Mansonella ozzardi*. *J Helminthol* 1977; 51:132-34.
6. Walker JS, Cadigan FC, Sirimanne AR, *et al.* Silvered leaf monkeys (*Presbytis cristatus*). I. Colony management, caging, diet and biologic values. *Lab Anim Sci* 1974; 24:290-98.
7. Palmieri JR, Van Dellen AF, Tirtokusumo S, *et al.* Trapping, care and laboratory management of the silvered leaf monkey (*Presbytis cristatus*). *Lab Anim Sci* 1984; 34:194-97.
8. Arambulo PV, Abass JB, Walker JS. Silvered leaf monkeys (*Presbytis cristatus*). II. Gastrointestinal parasites and their treatment. *Lab Anim Sci* 1974; 24:299-306.
9. Relethford JH, Coelho AM, Lawrence WA. Age estimation from dental eruption in infant and juvenile baboons (*Papio* sp). *Am J Primatol* 1982; 2:205-09.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Distribution of this document is unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NAMRU-2-TR-1044			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Naval Medical Research Unit No. 2		6b. OFFICE SYMBOL (If applicable) NAMRU-2	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) APO San Francisco, California 96528-5000			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Medical Research & Development Command		8b. OFFICE SYMBOL (If applicable) NMRDC	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62770A	PROJECT NO. 3M162770A870	TASK NO. AN
					WORK UNIT ACCESSION NO. 314
11. TITLE (Include Security Classification) (U) The silvered leaf monkey (<u>Presbytis cristata</u>) as a model for human bancroftian filariasis					
12. PERSONAL AUTHOR(S) J.R. Campbell; Harijani A. Marwoto*; S. Tirtokusumo; S. Masbar; J.T. Rusch Purnomo; B. Trenggono. *Indonesian Ministry of Health Research and Development					
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM 1987 TO		14. DATE OF REPORT (Year, Month, Day) 1987 August	
				15. PAGE COUNT 3	
16. SUPPLEMENTARY NOTATION Published in the Laboratory Animal Science, 37(4):502-504, August 1987. Done in cooperation with the Indonesian Ministry of Health.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	<u>Presbytis cristata</u> <u>Bancroftian filariasis</u>		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The silvered leaf monkey, <u>Presbytis cristata</u>, is presented as an experimental animal model for human bancroftian filariasis. In this paper baseline physical, hematological and parasitological data were obtained from 169 wild caught <u>P. cristata</u>. Animals were found naturally infected with a variety of protozoan and helminthic parasites including human pathogens. Bacteriological findings at time of capture were minimal although with increased human contact, gastrointestinal, respiratory and other infections among the animals became a significant management problem. <i>Reprinted from: Epidemiology, Reports 10. (1987)</i></p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL RAM I: Groves, M.			22b. TELEPHONE (Include Area Code) 301/663-7567		22c. OFFICE SYMBOL NMRDC